

1) Evaluate the line integral $\oint_C xy \, dx + x^2 y^3 \, dy$ where C is the triangle with vertices $(0, 0)$, $(1, 0)$, $(1, 2)$ by two methods:

a) Directly.

$$\frac{2}{3}$$

b) Using Green's Theorem.

$$\frac{2}{3}$$

Use Green's Theorem to evaluate the line integral along the given positively oriented curve.

2) $\int_C e^y \, dx + 2xe^y \, dy$ where C is the square with sides $x = 0$, $x = 1$, $y = 0$, and $y = 1$.

$$e - 1$$

3) $\int_C (y + e^{\sqrt{x}}) \, dx + (2x + \cos y^2) \, dy$ where C is the boundary of the region enclosed by the parabolas $y = x^2$ and $x = y^2$.

$$\frac{1}{3}$$

4) $\int_C xe^{-2x} dx + (x^4 + 2x^2y^2) dy$ where C is the boundary of the region between the circles $x^2 + y^2 = 1$ and $x^2 + y^2 = 4$.

0

5) $\int_C 2 \arctan \frac{y}{x} dx + \ln(x^2 + y^2) dy$ where $C: x = 4 + 2 \cos \theta, y = 4 + \sin \theta$.

0

Use Green's Theorem to evaluate $\int_C \vec{F} \cdot d\vec{r}$. (Check the orientation of the curve before applying the theorem.)

6) $\vec{F}(x, y) = \langle y^2 \cos x, x^2 + 2y \sin x \rangle$ where C is the triangle from $(0, 0)$ to $(2, 6)$ to $(2, 0)$ to $(0, 0)$.

-16

- 7) $\vec{F}(x, y) = (3x^2 + y)\mathbf{i} + 4xy^2\mathbf{j}$ where C is the boundary of the region lying between the graphs of $y = \sqrt{x}$, $y = 0$ and $x = 9$.

$$\frac{558}{5}$$

Use a line integral to find the area of the region R .

- 8) Region bounded by the graphs of $x^2 + y^2 = a^2$

$$\pi a^2$$

- 9) Triangle bounded by the graphs of $x = 0$, $3x - 2y = 0$ and $x + 2y = 8$.

$$4$$

- 10) Region bounded by the graphs of $y = 5x - 3$ and $y = x^2 + 1$.

$$\frac{9}{2}$$